

National Priority Chemicals Trends Report (2000-2004)

Section 4 Chemical Specific Trends Analyses for Priority Chemicals (2000–2004): Hexachloroethane (HCE)

Hazardous Waste Minimization and Management Division
Office of Solid Waste
U.S. Environmental Protection Agency

Contact Information:

Bill Kline, Senior Data Analyst Analysis & Information Branch (540) 341-3631 kline.bill@epa.gov

Tammie Owen, Data Analyst Analysis & Information Branch (703) 308-4044 owen.tammie@epa.gov

Ben Lesser, Chief Analysis & Information Branch (703) 308-0314 lesser.ben@epa.gov

Hexachloroethane (HCE)

Chemical Information:

CAS Number – 67–72–1

Alternate Names – carbon hexachloride, ethane hexachloride, perchloroethane

General Uses – HCE is mostly used by the military to make weapons that produce smoke, such as smoke pots and grenades used during training. It is also present as an ingredient in fungicides, insecticides, lubricants and plastics.

Potential Hazards – HCE is highly toxic; it may be fatal if inhaled, swallowed or absorbed through the skin.

Summary Analysis:

- NATIONAL: In 2004, approximately 3.8 million pounds of HCE were reported by nine facilities; this was approximately a 34 percent decrease compared to the quantity reported in 2000. Since 2000, the quantity of HCE decreased until it increased by approximately 1.1 million pounds or 40 percent in 2004. One facility accounted for 80 percent of the total quantity.
- REGIONAL: Facilities in four EPA regions where facilities reported HCE in 2004; since 2000, most of the HCE was reported by facilities in Region 6, including approximately 99 percent of the total quantity in 2004.
- STATES: Facilities in Louisiana reported approximately 89 percent of the total quantity of HCE in 2004, including an increase of approximately 1.4 million pounds in 2004. Most of this increase was reported by a facility in Louisiana which attributed much of the increase to improved flow measurement equipment.
- MANAGEMENT: Since 2000, treatment has been the primary method used to manage HCE; in 2004 approximately 96 percent of this chemical was treated.
- INDUSTRY SECTORS: Facilities in SIC 2869 (Industrial organic chemicals, nec) and SIC 2812 (Alkalies and chlorine) reported approximately 95 percent of the total quantity of HCE in 2004.

National Trends:

Exhibit 4.117 shows the number of facilities that reported HCE in 2000 to 2004 and the quantities of this PC that were managed via disposal, treatment, energy recovery, and recycling. In 2004, approximately 3.8 million pounds of HCE were reported by nine facilities. Compared to the quantity reported in 2000, there was approximately a 34 percent decrease in 2004. Since 2000, the quantity of HCE showed a trend of decreasing until 2004, when the quantity increased by approximately 1.1 million pounds, or 40 percent.

Since 2000, treatment has been the primary method used to manage HCE; in 2004 approximately 96 percent of this chemical was treated. Significant quantities of HCE were recycled each year, including approximately 2.3 million pounds in 2003 and in 2004.

Exhibit 4.117. National Management Methods for Hexachloroethane, 2000–2004

| Management Methods for HCE and Number of Facilities | 2000 | 2001 | 2002 | 2003 | 2004 | Percent Change (2000-2004) | Management Method Percent of Quantity of This PC (2004) |
|---|-----------|-----------|-----------|-----------|-----------|----------------------------------|--|
| Number of Facilities | 9 | 8 | 9 | 9 | 9 | 0.0% | - |
| Disposal Quantity (pounds) | 2,482 | 233 | 306 | 254 | 263 | -89.4% | 0.0% |
| Energy Recovery Quantity (pounds) | 1,245,190 | 455,985 | 206,953 | 139,929 | 146,243 | -88.3% | 3.9% |
| Treatment Quantity (pounds) | 4,462,309 | 3,689,031 | 3,849,238 | 2,553,948 | 3,626,347 | -18.7% | 96.1% |
| Priority Chemical Quantity (pounds) | 5,709,981 | 4,145,249 | 4,056,497 | 2,694,131 | 3,772,853 | -33.9% | - |
| Recycling Quantity (pounds)* | 1,027,963 | 850,000 | 3,530,419 | 2,336,505 | 2,279,804 | 121.8% | - |

^{*}Note: Waste minimization is the emphasis of this Report. As such, we primarily focus on quantities of PCs that are managed via onsite/offsite disposal, treatment, or energy recovery because we believe these PC quantities offer the greatest opportunities for waste minimization. Because recycled quantities of PCs are already directed to their best uses, they are considered separate and distinct from the quantities of PCs not recycled. Throughout this section, the recycled quantity is presented to provide some perspective regarding the quantity of this PC already recycled compared to the quantities that are managed via disposal, treatment, and energy recovery and thus potentially available for waste minimization

Exhibit 4.118 shows the number of facilities that reported HCE within various quantity ranges. Of the nine facilities that reported HCE in 2004, one facility accounted for 80 percent of the total quantity and five facilities reported approximately 97 percent of the total quantity of this chemical.

Exhibit 4.118. Distribution of Quantities by Facilities Reporting Hexachloroethane, 2004

| HCE (3,772,853 pounds) | | | | | | | | | |
|----------------------------|--|--|--|--|--|--|--|--|--|
| Quantity Reported | Number of Facilities Reporting This Quantity (2004) | Percent of Total Quantity of This PC (2004) | | | | | | | |
| up to 10 pounds | 0 | 0.0% | | | | | | | |
| 11 - 100 pounds | 0 | 0.0% | | | | | | | |
| 101 - 1,000 pounds | 1 | less than 0.1% | | | | | | | |
| 1,001 - 10,000 pounds | 0 | 0.0% | | | | | | | |
| 10,001 - 100,000 pounds | 3 | 2.7% | | | | | | | |
| 100,001 - 1 million pounds | 4 | 17.9% | | | | | | | |
| > 1 million pounds | 1 | 79.4% | | | | | | | |

EPA Regional Trends:

Exhibits 4.119 and 4.120 show the quantity of HCE for those five EPA regions where facilities reported this PC in 2000–2004; in 2004 facilities in only four regions reported HCE. Since 2000, most of the HCE was reported by facilities in Region 6, including approximately 99 percent of the total quantity in 2004. From 2000 to 2003 the quantity of HCE steadily decreased; however, in 2004 the quantity increased by approximately 1.1 million pounds, or 40 percent. Most of this increase was reported by a facility in Louisiana which attributed much of the increase to improved flow measurement equipment. A facility located in Oregon (Region 10) that only began reporting HCE in 2003 reported a decrease of approximately 73,000 pounds or 80 percent in 2004.

Exhibit 4.119. Regional Quantities of Hexachloroethane, 2000-2004

| EPA Region | 2000 (pounds) | 2001 (pounds) | 2002 (pounds) | 2003 (pounds) | 2004 (pounds) | Percent Change in Quantity (2000-2004) | Percent of Total Quantity of This PC (2004) |
|---------------|------------------|------------------|------------------|------------------|------------------|---|---|
| 4 | 0 | 0 | 84,900 | 0 | 0 | NA | 0.0% |
| 5 | 70,764 | 63,652 | 63,076 | 11,549 | 11,529 | -83.7% | 0.3% |
| 6 | 5,638,985 | 4,081,334 | 3,907,982 | 2,590,994 | 3,742,765 | -33.6% | 99.2% |
| 7 | 232 | 263 | 539 | 333 | 693 | 198.7% | 0.0% |
| 10 | 0 | 0 | 0 | 91,255 | 17,867 | NA | 0.5% |
| Total | 5,709,981 | 4,145,249 | 4,056,497 | 2,694,131 | 3,772,853 | -33.9% | 100.0% |

Exhibit 4.120. Distribution of Facilities Reporting Hexachloroethane in 2004 and the Quantities of Hexachloroethane Reported in 2004, by EPA Region




Exhibit 4.121 shows how HCE was managed by facilities in the four EPA regions in 2004. In 2004, approximately 99 percent of the HCE was treated, mostly onsite — by facilities in Region 6. Facilities in Regions 7 and 10 also primarily treated HCE. One facility in Region 5 used offsite energy recovery for all of its HCE. A relatively small quantity of HCE was land disposed. HCE was recycled by facilities in Regions 5 and 6; facilities in each of these two regions recycled a significant quantity compared to the quantity otherwise managed via land disposal, treatment, or energy recovery.

State Trends:

Exhibit 4.122 shows the quantity of HCE reported by facilities in seven states from 2000 to 2004. Facilities in Louisiana accounted for approximately 89 percent of the total quantity of this chemical in 2004, including an increase of approximately 1.4 million pounds in 2004. Most of this increase was reported by a facility in Louisiana which attributed much of the increase to improved flow measurement equipment.

Although Texas facilities reported approximately 11 percent of the total quantity in 2004, they reported a decrease of approximately 2.8 million pounds, or 87 percent, compared to the quantity reported in 2000. Likewise, a facility in Oregon reported a significant decrease from the quantity it reported in 2003. A facility in Michigan reduced its HCE by approximately 83 percent since 2000.

Exhibit 4.121. Regional Management Methods for Hexachloroethane, 2004

| RAMIAN | (nolinge) of | Percent of HCE (2004) | Disposal (pounds) | | Energy R (pour | - | Treatr (pour | | Recycling (pounds) | |
|--------|--------------|--------------------------|--------------------|---------------------|------------------------------|-------------------------------|---------------------|----------------------|---------------------|----------------------|
| | | | Onsite Disposal | Offsite Disposal | Onsite Energy Recovery | Offsite Energy Recovery | Onsite Treatment | Offsite Treatment | Onsite Recycling | Offsite Recycling |
| 6 | 3,742,765 | 99.2% | 30 | 3 | 134,714 | 0 | 3,534,174 | 73,844 | 2,237,150 | 0 |
| 10 | 17,867 | 0.5% | 0 | 4 | 0 | 0 | 12,813 | 5,050 | 0 | 0 |
| 5 | 11,529 | 0.3% | 0 | 0 | 0 | 11,529 | 0 | 0 | 0 | 42,654 |
| 7 | 693 | 0.0% | 227 | 0 | 0 | 0 | 0 | 466 | 0 | 0 |
| Total | 3,772,853 | 100.0% | 257 | 7 | 134,714 | 11,529 | 3,546,987 | 79,360 | 2,237,150 | 42,654 |

Exhibit 4.122. State Quantity Trends for Facilities Reporting Hexachloroethane, 2000–2004

| | | Total Quantity | (pounds) of Hexa | achloroethane | | | | | |
|-------|-----------|----------------|------------------|---------------|-----------|--------------------------------------|--|---|--|
| State | 2000 | 2001 | 2002 | 2003 | 2004 | Change in Quantity (2000–2004) | Percent Change in Quantity (2000–2004) | Percent of Total Quantity of This PC (2004) | |
| LA | 2,454,853 | 1,506,255 | 918,178 | 1,945,275 | 3,342,716 | 887,863 | 36.2% | 88.6% | |
| TX | 3,184,132 | 2,575,079 | 2,989,804 | 645,719 | 400,049 | -2,784,083 | -87.4% | 10.6% | |
| OR | 0 | 0 | 0 | 91,255 | 17,867 | 17,867 | NA | 0.5% | |
| MI | 68,464 | 63,652 | 63,076 | 11,549 | 11,529 | -56,935 | -83.2% | 0.3% | |
| KS | 232 | 263 | 539 | 333 | 693 | 461 | 198.7% | 0.0% | |
| AL | 0 | 0 | 84,900 | 0 | 0 | 0 | NA | 0.0% | |
| IN | 2,300 | 0 | 0 | 0 | 0 | -2,300 | -100.0% | 0.0% | |
| Total | 5,709,981 | 4,145,249 | 4,056,497 | 2,694,131 | 3,772,853 | -1,937,128 | -33.9% | 100.0% | |

Exhibits 4.123 through 4.126 show the trends for the quantities of HCE in the top five states in which facilities reported this PC in 2004.




Exhibit 4.124. Oregon Trends for Hexachloroethane, 2000–2004





Exhibit 4.125. Michigan Trends for Hexachloroethane, 2000-2004






Exhibit 4.126. Kansas Trends for Hexachloroethane, 2000-2004

Exhibits 4.127 and 4.128 show how HCE was managed by facilities in the two states in which facilities reported approximately 99 percent of the total quantity of HCE in 2004. Facilities in both these states primarily used onsite treatment to manage the HCE and reported similar recycled quantities.

Exhibit 4.127. Management Methods for Hexachloroethane, Facilities in States With 99 Percent of the Total Quantity, 2004

| State | Total Quantity (pounds) of HCE (2004) | Onsite Disposal (pounds) | Offsite Disposal (pounds) | Onsite Energy Recovery (pounds) | Offsite Energy Recovery (pounds) | Onsite Treatment (pounds) | Offsite Treatment (pounds) | Onsite Recycling (pounds) | Offsite Recycling (pounds) |
|-------|---|--------------------------------|---------------------------------|--|---|---------------------------------|----------------------------------|---------------------------------|----------------------------------|
| LA | 3,342,716 | 1 | 0 | 134,714 | 0 | 3,206,938 | 1,063 | 1,100,000 | 0 |
| TX | 400,049 | 29 | 3 | 0 | 0 | 327,236 | 72,781 | 1,137,150 | 0 |




Exhibit 4.128. State Distribution of Management Methods of Hexachloroethane, 2004

Industry Sector (SIC) Trends:

Exhibit 4.129 shows the quantity of HCE for the six industry sectors where facilities reported this chemical in 2000–2004.

- Facilities in SIC 2869 (Industrial organic chemicals, nec) and SIC 2812 (Alkalies and chlorine) reported approximately 95 percent of the total quantity of HCE in 2004.
- Since 2002, the quantity of HCE reported by facilities in SIC 2869 increased by approximately 1.1 million pounds. Part of this increased quantity was due to a change in the primary SIC code (from SIC 2812 to SIC 2869) by one of the larger facilities reporting HCE in 2002; the same facility also attributed much of the large increase in 2004 to improved flow measurement equipment at its facility. Conversely, this SIC change caused the SIC 2812 quantity to decrease in 2002. The quantity for SIC 2812 also decreased due to a large decrease in the reported quantity by one of the larger facilities in this industry sector and that facility's subsequent change in primary SIC code (from SIC 2812 to SIC 2821) in 2004. This SIC change, in turn, caused a significant increase in the quantity reported by SIC 2821 facilities in 2004.
- A facility in SIC 9711 (National security) that only began reporting HCE in 2003 reported a decrease of approximately 73,000 pounds or 80 percent in 2004. This facility reduced its use of HCE as a "surrogate" in trial burns done to ensure proper operation of its incinerators.

Exhibit 4.129. Industry Sectors in Which Facilities Reported Hexachloroethane, 2000-2004

| Primary SIC | SIC Description | Number of Facilities That Reported HCE (2004) | 2000 (pounds) | 2001 (pounds) | 2002 (pounds) | 2003 (pounds) | 2004 (pounds) | Change in Quantity (2000-2004) | Percent of Total Quantity of This PC (2004) |
|----------------|-----------------------------------|--|------------------|------------------|------------------|------------------|------------------|--------------------------------------|--|
| 2869 | Industrial organic chemicals, nec | 3 | 483,682 | 266,064 | 801,027 | 1,900,386 | 3,229,682 | 2,746,000 | 85.6% |
| 2812 | Alkalies and chlorine | 3 | 5,155,535 | 3,815,533 | 3,107,494 | 690,941 | 346,543 | -4,808,992 | 9.2% |
| 2821 | Plastics materials and resins | 2 | 68,464 | 63,652 | 63,076 | 11,549 | 178,762 | 110,298 | 4.7% |
| 9711 | National security | 1 | 0 | 0 | 0 | 91,255 | 17,867 | 17,867 | 0.5% |
| 3365 | Aluminum foundries | 0 | 2,300 | 0 | 0 | 0 | 0 | -2,300 | 0.0% |
| 3795 | Tanks and tank components | 0 | 0 | 0 | 84,900 | 0 | 0 | 0 | 0.0% |
| | Total | 9 | 5,709,981 | 4,145,249 | 4,056,497 | 2,694,131 | 3,772,853 | -1,937,128 | 100.0% |

Exhibit 4.130 shows how HCE was managed by the nine facilities in the four industry sectors that reported this chemical in 2004. Facilities in these four industry sectors used onsite treatment for approximately 94 percent of the HCE. Facilities in SIC 2812 (Alkalies and chlorine) used onsite energy recovery for approximately 27 percent of their HCE. Only approximately 500 pounds of HCE were land disposed. Two facilities, one each in SIC codes 2812 and 2821, reported approximately 98 percent of 2.2 million pounds of HCE recycled in 2004.

Exhibit 4.130. Management Methods for Hexachloroethane in Industry Sectors, 2004

| | | Total | Total Percent | | Disposal | | Energy Recovery | | ment | Recycling | |
|----------------|--------------------|------------------------------|--------------------------------|--------------------------------|---------------------------------|--|---|---------------------------------|----------------------------------|---------------------------------|----------------------------------|
| Primary SIC | SIC Description | Quantity of HCE (2004) | of Total Quantity (2004) | Onsite Disposal (pounds) | Offsite Disposal (pounds) | Onsite Energy Recovery (pounds) | Offsite Energy Recovery (pounds) | Onsite Treatment (pounds) | Offsite Treatment (pounds) | Onsite Recycling (pounds) | Offsite Recycling (pounds) |
| | Industrial organic | | | _ | _ | | _ | | | _ | _ |
| 2869 | chemicals, nec | 3,229,682 | 85.6% | 0 | 3 | 40,885 | 0 | 3,117,360 | 71,434 | 0 | 0 |
| | Alkalies and | | | | | | | | | | |
| 2812 | chlorine | 346,543 | 9.2% | 228 | 0 | 93,829 | 0 | 250,957 | 1,529 | 1,100,000 | 0 |
| | Plastics materials | | | | | | | | | | |
| 2821 | and resins | 178,762 | 4.7% | 29 | 0 | 0 | 11,529 | 165,857 | 1,347 | 1,137,150 | 42,654 |
| 9711 | National security | 17,867 | 0.5% | 0 | 4 | 0 | 0 | 12,813 | 5,050 | 0 | 0 |
| | Total | 3,772,853 | 100.0% | 257 | 7 | 134,714 | 11,529 | 3,546,987 | 79,360 | 2,237,150 | 42,654 |